

A. Shilf 10

LIBRARY

OF THE

U. S. Department of Agriculture.

Class

411

8-159

Sh 2

v. 19.

55343

HONORABILI ADMODUM
GULIELMO SCOTT,
EQUITI AURATO, &c. &c. &c.

VIRO OPTIMO ET LITERATISSIMO,

HUNC

DECIMUM NONUM
NATURÆ VIVARII
FASCICULUM

D. D. D.

GEORGIUS SHAW,

E. NODDER.

TO
THE RIGHT HONOURABLE
SIR WILLIAM SCOTT,
A GENTLEMAN
OF
DISTINGUISHED WORTH AND LITERATURE,
THIS NINETEENTH VOLUME
OF THE
NATURALIST'S MISCELLANY

IS
RESPECTFULLY INSCRIBED

BY
GEORGE SHAW,
E. NODDER.

LIBRARY
U.S. DEPARTMENT
OF AGRICULTURE

CORACIAS VARIA.

CHARACTER GENERICUS.

Rostrum cultratum, apice incurvato, basi pen-
nis denudatum.

Lingua cartilaginea, bifida.

Pedes ambulatorii.

Lin. Syst. Nat. p. 159.

CHARACTER SPECIFICUS.

Coracias albo nigroque varia, rostro plumbeo.

Coracias varia. *C. nigra*, alis albo variegatis,
dorso postico uropygio corporeque subtus
albis, rectricibus lateralibus albo terminatis.

Lath. ind. orn. p. 173.

Rarissimam hanc avem primus descripsisse videtur Buffonus, qui natale solum ei statuit novam Guineam. Ostenditur in tabula minor longe magnitudine naturali, quæ nempe pedalis est, addita quasi uncia. Quinam sint mores, et cui vivendi modo præcipue assuescat avis, incompertum est.

THE
PIED ROLLER.

GENERIC CHARACTER.

Bill strait, bending a little towards the end;
cultrated at the edges,

Nostrils narrow, naked.

Legs generally short: toes divided to their origin; three forwards and one backwards.

SPECIFIC CHARACTER.

Black-and-White Roller, with lead-coloured beak.

Cassican de la nouvelle Guinée.

Buff. ois. 7. p. 137.

Pl. Enl. 628.

Pied Roller.

Lath. Syn. 1. p. 415.

This very rare bird is, according to Buffon, by whom it appears to have been first described, a native of New Guinea. The plate represents it much reduced in size, its total length being about thirteen inches. Nothing particular seems to be known relative to its manners or history.



CEPHALUS BREVIS.

CHARACTER GENERICUS.

Maxillæ osseæ.

Corpus ovatum, postice truncatum, caput piscis simulans.

CHARACTER SPECIFICUS.

Cephalus brevis. C. corpore suborbiculato.

Tetrodon Mola. T. lævis compressus, cauda truncata, pinna brevissima, dorsali analique annexa.

Lin. Syst. Nat.

Diodon Mola. D. corpore lato, cauda truncata.

Bloch ichth. 4. t. 128.

Pisci huic insolita facie, quem ad genus *Tetrodontis* retulerat Linnæus, Blochiusque ad *Diodontis*, in opere nostro zoologico institutum est novum genus quod dicitur *Cephalus*. Maria incolit Europæa, corpore insignis paululum a capite protenso, et quasi derepente abrupto; adeo ut piscis cujuspian magni truncatum caput, magis quam animal absolutum et perfectum putes. In magnam crescere solet molem, diciturque in maribus nonnullis septentrionalibus in longitudinem extendi octopedalem, vel etiam decempedalem; et pondere æquare quingentas libras. Color plerumque fuscus, abdomine et lateribus argentatis. Comedere dicitur præcipue conchyliæ, et noctu lucem spargere splendide phosphoream.

THE
SHORT SUN-FISH.

GENERIC CHARACTER.

Jaws bony.

Body terminating abruptly, so as to resemble the head of a fish.

SPECIFIC CHARACTER.

Sun-Fish with suborbicular body.

Gen. Zool. 5. p. 437.


Sun-Fish.

Will. ichth. p. 151.

Short Diodon.

Penn. Brit. Zool.

This highly singular fish, which by Linnæus was considered as a species of *Tetrodon*, and by Dr. Bloch as a species of *Diodon*, was by myself stationed, in the work entitled *General Zoology*, under a new genus named *Cephalus*. It is a native of the European seas, and is remarkable for the very abrupt shape of its body, which terminating suddenly, at no great distance from the head, gives it rather the appearance of the truncated head of some large fish, than of an animal in its complete state. This fish grows to a very great size, and in some parts of the northern seas is said to have been sometimes seen of the length of eight or ten feet, and of the weight of five hundred pounds. Its general colour is brown, with a silvery cast on the sides and abdomen. It is supposed to feed principally on shell-fish, and is said to exhibit during the night a high degree of phosphoric splendor.



PAPILIO ADIPPE.

CHARACTER GENERICUS.

Antennæ apicem versus crassiores, sæpius clavato-capitatæ.

Alæ (sedentis) erectæ sursumque conniventes. (Volatu diurno).

Lin. Syst. Nat. p. 744.

CHARACTER SPECIFICUS.

Papilio Adippe. P. alis dentatis luteis nigro maculatis, subtus maculis 23? argenteis.

Lin. Syst. Nat. p. 786.

Nymph. phaler.

Papilio Adippe.

Fuessly Arch. ins. p. 1. pl. 1.

Inter insecta Britannica rariora numeratur Papilio Adippe, cujus larva violarum folia præcipue depascitur, et in chrysalidem mense Maio convertitur, e qua mense Junio erumpit papilio.



ADIPPE.

GENERIC CHARACTER.

Antennæ thickening towards the end, and commonly terminating in a club-shaped tip.

Wings (when at rest) meeting upwards. (Flight diurnal).

SPECIFIC CHARACTER.

Butterfly with denticulated luteous wings spotted with black, and marked beneath by twenty-three? silvery spots.

Moyen Papillon nacré.

Degeer ins. 2. p. 193.

pl. 1. f. 8. 9.

High brown Fritillary?

Harris Cat. Engl.

The *Papilio Adippe* is numbered among the rarer British insects: the caterpillar, which feeds principally on the leaves of violets, changes to a chrysalis in the month of May, and gives birth to the complete insect in June.



THE
LONG-FRONTED LOCUST.

GENERIC CHARACTER.

Head inflected, armed with jaws, and furnished with feelers.

Antennæ (in most species) setaceous or filiform.

Wings four, deflected and convoluted; the lower ones pleated.

Hind-Legs formed for leaping: claws on all the feet double.

SPECIFIC CHARACTER.

Green Locust, with conical head, and sword-shaped antennæ.

Long-headed African Locust.

The *Gryllus nasutus* or Long-fronted Locust, is a native of various parts of Africa, and is represented on the plate in its natural size.

COLUMBA AROMATICA.

CHARACTER GENERICUS.

Rostrum rectum, versus apicem descendens.

Nares oblongæ, membrana molli tumida semitectæ.

Lingua integra.

CHARACTER SPECIFICUS, &c.

Columba viridi-olivacea, alis nigris, fascia duplici flava.

Columba aromatica. *C. viridi-olivacea*, dorso castaneo, fascia alarum duplici flavicante et nigra, remigibus nigris, margine flavis.

Lath. ind. orn. p. 599.

In insulis Moluccis præcipue conspicitur *Columba aromatica*, minor paulo *columba* communi seu domestica. Insignitur avis alis nigrantibus, marginatis fascia duplici splendide flava alarum tectricibus tam minoribus quam majoribus.

COLUMBA AROMATICA.

CHARACTER GENERALIS.

Radix est, velut species herbarum.

Radix est, velut species herbarum. Radix est, velut species herbarum.

Radix.

Radix est, velut species herbarum.

CHARACTER SPECIFICUS.

Columba aromaticum, radix nigra, fascis du-

plici fascis.

Columba aromaticum, radix nigra, fascis du-

plici fascis, radix nigra, fascis du-

plici fascis, radix nigra, fascis du-

Radix est, velut species herbarum.

In hunc modum est, velut species herbarum. In hunc modum est, velut species herbarum.



THE
AROMATIC PIGEON.

GENERIC CHARACTER.

Bill strait, descending towards the tip.

Nostrils oblong, half covered by a soft tumid membrane.

Tongue entire.

SPECIFIC CHARACTER, &c.

Olive-green Pigeon, with black wings marked by a double yellow bar.

Pigeon vert d'Amboine.

Buff. ois. 2. p. 528.

Pl. Enl. 163.

Aromatic Pigeon.

Lath. syn. 2. p. 631.

The aromatic or Amboina Pigeon is found principally in the Molucca islands, and is of somewhat smaller size than the common or domestic pigeon. It is remarkable for the unusual colour of its wings, which are black, ornamented by a double bar of bright yellow; the smaller and larger coverts being tipped with that colour.

LABRUS TRIMACULATUS.

CHARACTER GENERICUS.

Dentes robusti, subacuti : molares interdum conferti, convexi : *labia* crassa, duplicata : *pinnæ dorsalis* radii, in certis speciebus, ramo elongati.

Opercula mutica, squamosa.

CHARACTER SPECIFICUS, &c.

Labrus ruber, squamis magnis, maculis tribus nigris ad finem dorsi.

Labrus trimaculatus. *L. ruber*, maculis duabus utrinque ad basin *pinnæ dorsalis*, tertiaque inter hanc et caudam.


Lin. Syst. Nat. Gmel. p. 1294.

Labrus carneus. *L. maculis* tribus nigris in parte posteriore dorsi.

Bloch ichth. 9. p. 1. t. 289.

Labrum trimaculatum marium Europæorum incolam, in litoribus etiam nostratibus interdum conspectum, depinxit Pennantus in opere suo zoologico Britannico. Crescit, ut plurimum, in longitudinem duodecim seu quindecim unciarum. Variat color, plus nempe seu minus saturatus; quod accidere solet et cæteris contribulibus.

706



22 Noddy, Vol. d. L. 1847

THE
TRIMACULATED LABRUS.

GENERIC CHARACTER.

Teeth strong and subacute: the grinders sometimes convex and crowded: *lips* thick and doubled: rays of the *dorsal fin*, in some species, elongated into soft processes.

Gill-Covers unarmed and scaly.

SPECIFIC CHARACTER, &c.

Red Labrus, with large scales, and three black spots towards the end of the back.

Gen. Zool. 4. p. 502.

Trimaculated Wrasse.

Penn. Brit. Zool. 3. p. 218.

Der rothe Lippfisch.

Bloch ichth. pl. 289.

This species, which is a native of the European seas, is occasionally observed about our own coasts, and is described by Mr. Pennant in the British Zoology. Its general length is about twelve or fifteen inches, and, like most others of its tribe, it is subject to considerable variation as to the intensity of its colour.

PAPILIO AMPHINOME.

CHARACTER GENERICUS.

Antennæ apicem versus crassiores, sæpius clavato-capitatae.

Alæ (sedentis) erectæ sursumque conniventes, (volatu diurno.)

Lin. Syst. Nat. p. 744.

CHARACTER SPECIFICUS, &c.

Papilio alis dentatis nigris cæruleo nebulosis, primoribus fascia utrinque alba, posticis (subtus) rubro radiatis.

Lin. Syst. Nat. p. 779

Nymph : phaler :

Mer. Surin. t. 7.

Roes. 4. t. 10. f. 1. 2.

Insolitæ pulchritudinis *Papilio* in tabula depingitur, Americæ australis præcipue Surinamiæ incola. Extat eleganter depictus in splendido opere Dominæ Merian de insectis Surinamensibus, quæ narrat larvam folia depasce Plumeriæ rubræ, et in chrysalidem mense junio converti, unde Julio insequente erumpit *papilio*.



AMPHINOME.

GENERIC CHARACTER.

Antennæ thickening towards the end, and commonly terminating in a clavated tip.

Wings (when at rest) meeting upwards.

Flight diurnal.

SPECIFIC CHARACTER, &c.

Butterfly with dentated black wings clouded with blue; the upper pair marked on both sides by a white band; the lower pair radiated beneath with red.

Aubent. Pl. Enl. 92. f. 7. 8.

Mer. ins. Surin. pl. 7.

Cramer ins. 5. pl. 54. E. F.

The highly elegant insect represented on the present plate is a native of South America, and more particularly of Surinam, where its caterpillar, according to Madam Merian, who has figured it in her splendid work on the insects of that country, feeds on the leaves of the *Plumeria rubra*, and changes into a chrysalis in the month of June, from which in the month of July emerges the complete insect.

GRYLLUS SQUARROSUS.

CHARACTER GENERICUS.

Caput inflexum, maxillosum, palpis instructum.

Antennæ setaceæ seu filiformes.

Alæ quatuor, deflexæ, convolutæ : inferiores plicatæ.

Pedes postici saltatorii : ungues ubique bini.

Lin. Syst. Nat. p. 692.

CHARACTER SPECIFICUS.

Gryllus thorace utrinque articulis tribus trispinosis, elytris viridibus, alis rubris, omnibus nigro punctatis.

Gryllus squarrosus. *G. viridis*, thorace subarticulato, articulis utrinque tricuspидibus, alis rubris nigro punctatis.

Lin. Mantiss. 553.

Gryllus squarrosus.

Fabr. sp. ins. 1. p. 364.

Drury exot. ins. 1. pl. 49. f. 1.

Varias Africæ regiones incolit *Gryllus squarrosus*, a Domino Drury in elegantissimo suo opere de insectis exoticis primum descriptus. Tabula insectum magnitudine vera exprimit.

GRYLUS AQUARIUS

CHARACTER GENERALIS

GRYLUS AQUARIUS

Caput inflexum, maxillae palpi instructum.

Antennae sexarticulae, sensu distinctae.

Alae quatuor, dorsales, convolutae; inferiores

plicatae.

Pede posteriori saltatori: ungues bipides bini.

Loc. nat. M. B. G. 2.

CHARACTER SPECIFICUS

Gryllus thorace utrinque articulis tribus tripli-

bus, elytris viridibus, alis rubris, omnino

hirsuto punctatis.

Gryllus aquaticus. C. viridis, thorace subar-

ticulato, articulis utrinque triplicibus,

alae rubris nigro punctatis.

Loc. nat. M. B. G. 2.

Gryllus aquaticus.

Tab. 1. p. 201.


Tab. 2. p. 10. 11.

Variet. Alae nigrae in G. gryllus aquaticus a

Donato Hym. in descriptionibus suis et in

extrema pagina descriptionis Tab. 1. p. 10. 11.

Aliae rubrae respicitur.





THE
TRIPLE-SPINED LOCUST.

GENERIC CHARACTER.

Head inflected, armed with jaws, and furnished with feelers.

Antennæ, in most species, setaceous or filiform.

Wings four, deflected and convoluted; the lower ones pleated.

Hind Legs formed for leaping: claws on all the feet double.

SPECIFIC CHARACTER, &c.

Locust with three triple-spined protuberances on each side the thorax; the elytra or upper-wings green, the lower red; all speckled with black.

The triple-spined African Locust.

The present species of Locust, which is represented on the plate in its natural size, is a native of several parts of Africa. It seems to have been first described in the elegant work of Mr. Drury on exotic insects.



THE
DUSKY PARROT.

GENERIC CHARACTER.

Bill hooked: upper mandible moveable, and furnished with a cere.
Nostrils in the base of the bill.
Tongue fleshy, obtuse, generally entire.
Feet scansorial.

SPECIFIC CHARACTER, &c.

Blackish Parrot, purplish beneath, with the wings and tail blue; the latter half-red beneath.

Little dusky Parrot.

Edw. p. 315.

Lath. ind. orn.

Papegai violet.

Buff. ois. 6. p. 244.

Perroquet varié de Cayenne.

Pl. Enl. 4

This species is a native of South America, and particularly of Surinam: its general size is that of a common domestic pigeon. In its colours it sometimes varies a little.

FISTULARIA CHINENSIS.

CHARACTER GENERICUS.

Rostrum cylindricum, apice maxillosum.

Corpus elongatum.

Membr: branch: radiis septem.

CHARACTER SPECIFICUS, &c.

Fistularia squamosa rufescens, nigro maculata,
cauda rotundata mutica.

Gen. Zool. 5. p. 97.

Fistularia Chinensis. F. cauda rotundata.

Lin. Syst. Nat. p. 515.

Fistularia edentata, cauda rotundata.

Bloch. ichth. t. 388.

In maribus Indicis præcipue conspicitur *Fistularia Chinensis*, in longitudinem tripedalem vel quadripedalem crescens, et vermibus insectisque marinis variis victitans.



THE
CHINESE FISTULARIA.

GENERIC CHARACTER.

Snout cylindric: mouth terminal.

Body lengthened.

Gill-Membrane seven-rayed.

SPECIFIC CHARACTER, &c.

Scaly, rufescent *Fistularia*, with black spots
and simple, rounded tail.

Gen. Zool. 5. p. 97.

The Chinese Trumpet-Fish.

This remarkable fish is principally seen in the Indian seas, where it grows to the length of three or four feet, and feeds on various kinds of marine worms and insects.

PAPILIO CASSIÆ.

CHARACTER GENERICUS.

Antennæ apicem versus crassiores, sæpius clavato-capitatae.

Alæ (seditis) erectæ sursumque conniventes, (volatu diurno.)

Lin. Syst. Nat. p. 744.

CHARACTER SPECIFICUS, &c.

Papilio Cassiæ. P. alis integerrimis fuscis; primoribus posticarumque margine fascia ferruginea, omnibus subtus ocellatis.

Lin. Syst. Nat. p. 767.

Dan : Festiv :

Mer. Surin. t. 32.

Surinamiam incolit hic Papilio, ubi larva ejus varias Cassiæ species depascitur, et in chrysalidem mense Maio convertitur, e qua mense Junio erumpit insectum plene formatum.



THE
CASSIA BUTTERFLY.

GENERIC CHARACTER.

Antennæ generally thickening towards the extremities, and terminating in a clavated tip.

Wings (when sitting) erect, and meeting upwards. Flight diurnal.

SPECIFIC CHARACTER, &c.

Butterfly with entire, brown wings ; the upper pair and the margin of the lower marked by a ferruginous band, and all marked beneath by eye-shaped spots.

Clerk ic. ins. t. 29. f. 3.

Cramer ins. g. pl. 106. A.

This species is a native of Surinam, where its caterpillar feeds on various species of Cassia. It changes to a chrysalis in the month of May, from which in June proceeds the complete insect.

APHRODITA CIRRHOSA.

CHARACTER GENERICUS.

Corpus repens oblongum, squamis tectum; pedibus utrinque setosis.

Os terminale, cylindricum.

Tentacula duo setacea annulata.

Oculi quatuor.

Lin. Syst. Nat. Gmel.

CHARACTER SPECIFICUS, &c.

Aphrodita cirrhosa. A. lineari-attenuata, pedunculis compressis apice bipartitis, squamis vesiculosus.

Lin. Syst. Nat. Gmel.


Aphrodita cirrhosa.

Pall. Misc. Zool. p. 95.

t. 8. f. 3.—6.

Hanc speciem ita describit celeberrimus Pallas, in opere cui titulus *Miscellanea Zoologica*.

“Corpus tenerrimum, longiusculum, lineari-attenuatum, ex tereti depressiusculum adeoque ad *Nereides* figura accedens. Pedunculi utrinque triginta-quinque, productiores, compressi, apice bipartiti, porzione superiore minore, utraque penicillo ex pilis fulvescentibus, in superiori longioribus, in inferiore numerosioribus barbata. Squamæ in dorso undecim parium, subrotundæ, areola in anteriore disci parte, supra pedunculos adnatæ margine postico scabriusculæ, fuscoque denticulatæ, in specimine quod descripsi, omnes in vesiculas inflatæ. Cuticula albida atque iridescens huic speciei, teneriorque est quam ulli congenerum.”



THE
CIRRHATED APHRODITA.

GENERIC CHARACTER.

Body repent, oblong, in most species covered with scales, and furnished on each side with bristle-shaped feet.

Mouth terminal, cylindric.

Feelers two, setaceous, annulated.

Eyes four.

SPECIFIC CHARACTER, &c.

Pale, linear Aphrodita, with eleven pair of large rounded, vesicular scales along the back, and compressed feet divided at the tip.

Large-Scaled Aphrodita.

This species is described in the work of Dr. Pallas, entitled *Miscellanea Zoologica*. The body, says Dr. Pallas, is very tender, lengthened, linear-attenuated, flattish-rounded, and consequently approaching to the form of the genus *Nereis*. On each side are thirty-five lengthened, compressed feet, divided at the tip, the superior portion smaller, and each bearded by a pencil of subfulvous hairs, longer on the upper portion, but more numerous on the lower. On the back are eleven pair of scales, which are roundish, situated above the feet, rather rough on their hinder margin, and marked with fuscous denticulations. In the specimen described all the scales were inflated into vesicles. The skin in this species is whitish and iridescent, and more tender than in any of its congeners.

ALCEDO LEUCOCEPHALA.

CHARACTER GENERICUS.

Rostrum trigonum, crassum, rectum, longum.

Lingua carnea, brevissima, plana, acuta.

Pedes gressorii plerisque.

Lin. Syst. Nat. p. 178.

CHARACTER SPECIFICUS, &c.

Alcedo viridi-cærulea, alis caudaque nigro-cæruleis, capite corporeque subtus stramineis, rostro rubro.

Alcedo leucocephala. *A.* ex cæruleo viridis, capite collo et subtus alba, remigibus fuscis.

Lin. Syst. Nat. Gmel.

Insulam Javam incolit formosa hæc avis, magnitudine fere duplo major *Alcedine Ispida* sive *Europæa*.



Alcedo alata

THE
WHITE-HEADED KINGFISHER.

GENERIC CHARACTER.

Bill trigonal, thick, strait, long.

Tongue fleshy, very short, flat, sharp-pointed.

Feet gressorial.

SPECIFIC CHARACTER, &c.

Blue-green Kingfisher, with blackish-blue wings
and tail; the head and under parts straw-
coloured.

Martin-Pêcheur de Java.

Pl. Enl. 757.

White-Headed Kingfisher.

Lath. Syn. p. 617.

This beautiful species is a native of the island of
Java, and is nearly twice the size of the common
European Kingfisher.

794



R. P. N. & Co. Del. & Sculp.

THE
CRETAN SCARUS.

GENERIC CHARACTER.

Jaws long, divided in the middle, crenated on the edge; the teeth connate and conglomerate.

SPECIFIC CHARACTER, &c.

Green Scarus, yellowish beneath, with very large scales, ramified lateral line, and sub-lunated tail.

Gen. Zool. p. 393.

Der Griechische Papageyfisch.

Bloch. ichth. 7. pl. 220.

The Cretan Scarus grows to the length of about twelve inches, and is found both in the European and Indian seas.

SPHINX SATELLITIA.

CHARACTER GENERICUS.

Antennæ medio crassiores, seu utraque extremitate attenuatæ, subprismaticæ.

Alæ deflexæ (volatu graviore vespertino seu matutino).

Lin. Syst. Nat. p. 796.

CHARACTER SPECIFICUS, &c.

Sphinx alis integris griseo-olivaceis fusco nebulosis, primoribus puncto duplici ocellato nigro.

Sphinx Satellitia. S. alis griseo nebulosis, primoribus puncto nigro ocellari sesquialtero, posterioribus basi cinereis.

Lin. Syst. Nat. Gmel.

Sphinx Satellitia.

Fabr. sp. ins. 2. p. 148.

Drury ins. 1. t. 29. f. 1. 2.

Jamaicam incolit Sphinx Satellitia, magnitudine vera in tabula depicta.

795



Sphinx morio

THE
DOUBLE-POINTED SPHINX.

GENERIC CHARACTER.

Antennæ subprismatic, thickest in the middle,
and attenuated at each extremity.

Wings deflected. Flight generally matutine or
vespertine.

SPECIFIC CHARACTER, &c.

Sphinx with entire olive-grey wings clouded
with brown, the upper pair marked by a
double ocellated black speck.

The Double-Pointed Hawk-Moth.

Double-Pointed Jamaica Sphinx.

This insect is a native of Jamaica, and is represented on the plate in its natural size.

BRACHIONUS ACULEATUS.

CHARACTER GENERICUS.

Corpus contractile, testa tectum, ciliis rotatoriis.

Lin. Syst. Nat. Gmel,

CHARACTER SPECIFICUS.

Brachionus testa oblongo-quadrata, antice quadricorni, postice bicorni, capite tentaculis setaceis divergentibus.

Detectum est mense Julio in aqua stagnante insigne animalculum quod aucta admodum naturali magnitudine in tabula depingitur. Motu fertur celerrimo, testa obtectum pellucida, fronte in laminas duas divisa, angulis spina incurva terminatis. Instruitur caput fasciculis pluribus vibrissarum seu setarum divergentium, quæ tamen, non nisi movente se animalculo, visibiles sunt. Cor, prope partem superiorem situm, cordique simillimum Vorticellæ quæ *rotatoria* dicitur, pulsu protruditur evidenti et celerrimo. Cætera viscera, non sigillatim cernenda, fusco-flavescent.

796



Rur

THE
ACULEATED BRACHIONUS.

GENERIC CHARACTER.

Body contractile, inclosed in a shell, and furnished at the head with rotatory organs.

SPECIFIC CHARACTER.

Brachionus with oblong-square shell with four spines in front, two behind, and head furnished with divergent setaceous tentacula.

The remarkable animalcule exhibited in a highly magnified state on the present plate, was discovered in stagnant water in the month of July. It is an animal of very swift motion, and appears covered with a transparent shell of an oblong square shape, divided into two laminæ in front, and terminated at each extremity by a curved spine. The head or upper part is furnished with several groupes or fasciculi of divergent vibrissæ or strait setaceous fibres, which are only apparent while the animal is in motion. The heart, which is situated near the upper part, is very apparent, exerts a very quick pulsation, and much resembles that of the *Vorticella rotatoria* or Wheel-Animal in shape. The other viscera are not separately distinguishable, and are of a yellowish-brown colour.



R. P. Nodder Del. et Sculp. t.

THE
PURPLE-THROATED CREEPER.

GENERIC CHARACTER.

Bill bent, slender, sharp-pointed.
Tongue differing in different species.
Feet formed for walking.
Tail consisting of twelve feathers.

SPECIFIC CHARACTER, &c.

Brown Creeper, with green front, and purple throat.

The Purple-Throated Creeper.

Brasilian green-faced Creeper.

Grimpereau du Bresil.

Pl. Enl. 578. f. 3.

The species here represented is a native of Brasil, and is figured in its natural size.

TUBULARIA RAMOSA.

CHARACTER GENERICUS.

Stirps tubulosa, simplex vel ramosa, basi affixa.

Animal terminale, capite tentaculis cristato.

CHARACTER SPECIFICUS, &c.

Tubularia culmis ramosis, geniculis contortis.

Lin. Syst. Nat. p. 1302.

Corallina tubularia gracilis et ramosa, axillis ramulorum contortis.

Ellis Corall. p. 31. t. 17. A. a.

In litoribus Europæis non raro reperitur hoc zoophyton, carens quidem magnitudine, non tamen elegantia. Adhæret conchis, rupibus, aliisque marinis. Depingitur in tabula tam naturalis moles, quam modice aucta.

TUBULARIA RAMOSA.

CHARACTERES GENERALES.

CHARACTERES SPECIFICI.

Superior tubulosa, simplex vel ramosa, breviter
Animal teretiuscula, capite tentaculis cincto.

CHARACTERES SPECIFICI.

Tubularia cylindrica, ramosa, geniculis contractis.
Tub. Gen. Nov. p. 130.
Corallium tubulosa geniculis et ramosa, axillis
ramulorum contractis.
J. M. Cuv. p. 31. t. 17. n. 2.

Le tubulaire Lamourin non est répété par les
autres auteurs qui ont mentionné, non tant le
genre. Adhuc quodammodo, répété, plusieurs fois.
Distingue le tubulaire non répété par les autres
des autres.



Scilla maritima L.

THE
TWISTED TUBULARIA.

GENERIC CHARACTER.

Stem tubular, simple or branched, fixed by the base.

Animal terminal; the head crested with tentacula.

SPECIFIC CHARACTER, &c.

Branched Tubularia, with the axillæ of the joints twisted.

Small ramified tubular Coralline.

Ellis Corall. p. 31. pl. 17.

This small but elegant zoophyte is a frequent inhabitant of the European coasts, adhering to rocks, shells, and various other marine substances. It is represented on the plate both in its natural size, and as it appears when moderately magnified.

CHÆTODON TEIRA.

CHARACTER GENERICUS.

Caput parvum. *Os* exiguum: *Dentes* conferti, flexiles, setacei.

Membr. branch. radiis tribus ad sex.

Corpus latum, compressum, pluribus fasciatum; pinna *dorsi anique* basi crassa, carnosa, squamosa.

CHARACTER SPECIFICUS, &c.

Chætodon albus, corpore elevato-orbiculari, pinna dorsali analique magnis falcatis longissimis.

Chætodon Teira. *C. fasciis* tribus nigris, pinna dorsali analique longissimis.

Bloch ichth. t. 199, f. 1.

Piscibus quos amplectitur genus *Chætodon* peculiaris quædam sæpius est conformatio: qua tamen in re pauci insigniores illo de quo jam agitur, in maribus Indicis generato; cujus moles interdum ampla esse dicitur, licet parva, ut plurimum, sint specimina quæ videre est in museis Europæis.



THE
LONG-FINNED CHÆTODON.

GENERIC CHARACTER.

Head small. *Mouth* small: *Teeth* close-set, flexile, setaceous.

Gill-Membrane three, four, five, or six-rayed.

Body broad, compressed, and generally fasciated: *dorsal* and *anal* fin thick, fleshy and scaly at the base.

SPECIFIC CHARACTER, &c.

White Chætodon with high-orbicular body, and large, extremely long, falcated dorsal and anal fin.

Long-Finned Chætodon.

Gen. Zool. 4. p. 345.

Many fishes of the genus Chætodon are remarkable for the singularity of their shape; few however will be found to exceed in this particular the species here represented, which is an inhabitant of the Indian seas, and is said to arrive at a considerable size, though the specimens usually seen in Museums are but small.

BUCCINUM SUBULATUM.

CHARACTER GENERICUS.

Animal Limax.

Testa univalvis, spiralis, gibbosa.

Apertura ovata, desinens in canaliculum dextrum, cauda retusum.

Labium interius explanatum.

Lin. Syst. Nat. p. 1198.

CHARACTER SPECIFICUS, &c.

Buccinum testa subulata lævi flava, maculis subquadratis duplicatis fuscis.

Buccinum subulatum. B. testa turrita subulata lævi indivisa integerrima.

Lyn. Syst. Nat.

Rumph. mus. t. 30. f. B.

Bonann. recr. t. 118.

In maribus Indicis invenitur Buccinum subulatum
cujus veram magnitudinem cernere est in tabula.



THE
SUBULATE BUCCINUM.

GENERIC CHARACTER.

Animal allied to a slug.

Shell univalve, spiral, gibbous.

Aperture ovate, ending in a channel pointing
towards the right.

Interior Lip expanded.

SPECIFIC CHARACTER, &c.

Buccinum with smooth yellow subulate shell,
marked by a double row of squarish brown
spots.

Smooth double-spotted Buccinum.

Yellow screw Buccinum.

This shell is an inhabitant of the Indian seas, and is
represented on the plate in its natural size.

THE
SUBULATE BUCCIUM.

GENERAL CHARACTER.

Animal allied to a dog.
Shell ovate, spiral, gibbous.
Aperture ovate, ending in a channel pointing
towards the right.
Lateral lip expanded.

TYPICAL CHARACTER.

Buccinum with smooth yellow subulate shell,
marked by a double row of squarish brown
spots.
Smooth double-spotted Buccinum.
Yellow screw Buccinum.

This shell is an imitation of the Indian seal, and is
represented on the plate in its natural size.

STRIX LARVATA.

CHARACTER GENERICUS.

Rostrum aduncum, absque cera.

Nares oblongæ, pennis setaceis recumbentibus
obtectæ.

Caput grande, auribus oculisque magnis.

Lingua bifida.

Digitus externus retro mobilis.

CHARACTER SPECIFICUS, &c.

Strix alba, alis subferrugineis, facie nigra.

Strix personata.

Daudin orn. 1. p. 192.

Avem depinximus elegantem, Cayanæ in America
australi incolam; quæque fortasse varietas esse possit
Strigis perspillatæ Latham. In majoribus habetur sui
generis, longa nempe quasi viginti uncias.



W. Woodcock Del. et Sculp.

THE
MASKED OWL.

GENERIC CHARACTER.

Bill hooked, without cere.

Nostrils oblong, covered with recumbent setaceous feathers.

Head, eyes, and ears, large.

Tongue bifid.

Exterior toe moveable backward.

SPECIFIC CHARACTER, &c.

White Owl, with subferruginous wings and black face.

La Chouette masquée.

Levaill. ois. pl. 44.

The elegant bird here represented is a native of Cayenne in South America, and is perhaps no other than a variety of the *Strix perspicillata* or Spectacle Owl of Mr. Latham. It is one of the larger birds of the genus, measuring about twenty inches in length.

ACANTHURUS CHIRURGUS.

CHARACTER GENERICUS.

Dentes parvi, plerisque lobati.

Cauda utrinque aculeata.

CHARACTER SPECIFICUS, &c.

Acanthurus luteus corpore postice striis transversis fuscis.

Acanthurus Chirurgus.

Gen. Zool. 4. p. 379.

Chætodon Chirurgus.

Bloch. ichth. t. 208.

In opere hoc nostro non ita pridem exposuimus quæ de causa certas Linnæani generis *Chætodontis* species, mutato nomine, *Acanthuburos* nuncupavimus. *Acanthuri* species quæ in tabula ostenditur in maribus nutritur Americanis, raro longior octo seu decem unciis.



THE
LANCET-TAILED ACANTHURUS.

GENERIC CHARACTER.

Teeth small, in most species lobated.

Tail aculeated on each side.

SPECIFIC CHARACTER, &c.

Orange-yellow Acanthurus, with the body
crossed on the hind-part by transverse
brown stripes.

Lancet Acanthurus.

General Zoology, vol. 4. p. 379.

The reason for placing certain species of the Lin-
næan genus *Chætodon* under a separate genus entitled
Acanthurus, has been already explained in the present
work. The species of *Acanthurus* represented on the
annexed plate is a native of the American seas, where
it is rarely observed to measure more than eight or ten
inches in length.

SPHINX POPULI.

CHARACTER GENERICUS.

Antennæ subprismaticæ, utroque fine attenuatæ.

Lingua exserta (plerisque).

Palpi duo reflexi.

Alæ deflexæ.

CHARACTER SPECIFICUS, &c.

Sphinx alis dentatis griseo-gilvis reversis, posticis basi ferrugineis.

Sphinx Populi. S. alis dentatis reversis, posticis basi ferrugineis, anticis puncto albo.

Lin. Syst. Nat.

Sepp ins. 3. p. 3. pl. 1.

Merian Ins. Europ. pl. 37.

Folia populi salicisque depascitur larva Sphingis hujus Britannicæ. In chrysalidem mense Augusto convertitur, e qua Maio vel Junio insequente erumpit ipsa Sphinx.



THE
POPLAR SPHINX.

GENERIC CHARACTER.

Antennæ subprismatic, attenuated at each extremity.

Tongue generally exerted.

Feelers two, reflex.

Wings deflected.

SPECIFIC CHARACTER, &c.

Sphinx with dentated, buff-grey, reversed wings; the lower pair ferruginous at the base.

The Poplar Hawk-Moth.

Degeer ins. 1. pl. 8. f. 5.

Roes. ins. 3. pl. 30.

The Caterpillar of this species, which is a native of our own country, feeds on the leaves of the willow and poplar, changing to a chrysalis in the month of August, out of which in May or June following, emerges the complete Sphinx.

PAPILIO ANDROGEUS?

CHARACTER GENERICUS.

Antennæ apicem versus crassiores, sæpius clavato-capitatæ.

Alæ (sedentis) erectæ sursumque conniventes. (Volatu diurno).

Lin. Syst. Nat. p. 744.

CHARACTER SPECIFICUS, &c.

Papilio alis dentatis nigro æneis, posticis subtus lunulis rubris cyaneis flavisque.

Fab. spec. ins. 2. p. 8.

Pap. Androgeus?

Cramer ins. 2. t. 16. f. c. d.

Pap. Polydamas?

Lin. Syst. Nat. 747.

Pap. Peranthus?

Cram. t. 204. A. B.

In America australi, præcipue in Surinamia, conspicitur Papilio Androgeus, cujus larva hibiscorum folia depascitur. In tabula nostra magnitudine vera una cum larva pupaque ostenditur.



ANDROGEUS.

GENERIC CHARACTER.

Antennæ commonly thickening towards the end into a clavated tip.

Wings (when at rest) meeting upwards. Flight diurnal.

SPECIFIC CHARACTER, &c.

Butterfly with blackish-green brassy wings, the lower pair marked beneath by a verge of red, blue, and yellow crescents.

Mer. Sur. pl. 31.

Kleeman ins. pl. 8.

The present species is a native of South America, and particularly of Surinam, where its caterpillar feeds on the leaves of the genus *Hibiscus*. The plate represents it in its different states and in its natural size.

STURNUS SUBROSEUS.

CHARACTER GENERICUS.

Rostrum subulatum, angulatum, depressum, obtusiusculum: mandibula superiore integerrima, marginibus patentiusculis.

Nares supra marginatæ.

Lingua emarginata, acuta.

Lath. ind. orn.

CHARACTER SPECIFICUS, &c.

Sturnus griseo-rufescens, subtus subroseus, vertice cristato remigibusque primoribus nigris.

Turdus Pagodarum?

Lath. ind. orn.

Le Martin brame.

Levaill. ois. t. 95.

Testatur Levaillant, hujus avis descriptor, *Sturnis* subroseis abundare multas Indicas et Africanas regiones; seque ipsum eos conspexisse gregatim collectos. Si habitum generalem spectes, simillima est *Turdó roseo*, nisi quod colores sint obscuriores. Ipsissima est magnitudine qua *Sturnus* vulgaris; quem inter et hanc avem propinquam esse cognationem suspicatur Levaillant.



THE
SUBROSACEOUS STARE.

GENERIC CHARACTER.

Bill subulate, angular, depressed, somewhat obtuse; the upper mandible entire.

Nostrils marginated above.

Tongue emarginate, sharp-pointed.

SPECIFIC CHARACTER, &c.

Rufescent-grey Stare, subrosaceous beneath, with black crested crown and black primary wing-feathers.

Le Martin brame.

Levaill. ois. pl. 95.

This bird, according to its describer Levaillant, is very common in many parts of India as well as of Africa, where considerable flights were observed by himself. In its general appearance it seems considerably allied to the *Turdus roseus*, or Rose-coloured Ouzel, but is of less vivid colours. Mons. Levaillant seems to consider it as approaching very nearly to the Starling tribe, and in size it is exactly equal to the common Starling.

PAPILIO SPONDIÆ.

CHARACTER GENERICUS.

Antennæ apicem versus crassiores, sæpius clavato-capitatae.

Alæ (sedentis) erectæ sursumque conniventes (volatu diurno).

Lin. Syst. Nat.

CHARACTER SPECIFICUS, &c.

Papilio alis denticulatis cæruleis, subtus fuscis, inferioribus macula marginali flava.

Mer. Sur. t. 13.

Eruca papilionis in tabula depicti, quam edacissimam esse asserit Domina Merian, folia depascitur arboris quæ *Spondias lutea* dicitur. In chrysalidem convertitur ineunte mense Aprili, quo fere exacto, erumpit insectum plene formatum. In Surinamia generatur Papilio Spondiæ; cujus vera magnitudo una cum eruca et chrysalide in tabula ostenditur.



THE
SINGLE-SPOTTED BUTTERFLY.

GENERIC CHARACTER.

Antennæ thickening towards the end, ending in a clavated tip.

Wings (when sitting), erect and meeting upward ; flight diurnal.

SPECIFIC CHARACTER, &c.

Blue Butterfly, with denticulated wings, brown beneath, with a yellow marginal spot on the lower pair.

Mer. Sur. pl. 13.

The caterpillar of the Butterfly here represented, and which, according to Madam Merian, is of a very voracious nature, feeds on the leaves of the *Spondias lutea*, or American Plum ; changing to chrysalis in the beginning of April, and producing the complete insect towards the end of the same month. The plate shews the animal in all its states, and in its natural size. It is a native of Surinam.

SPARUS FALCATUS,

CHARACTER GENERICUS.

Dentes validi; Primores in aliis ordine simplici, in aliis duplici, triplici, vel quadruplici dispositi.

Molares (plerisque) validi, convexi, læves, in series dispositi, et quasi pavementum in ore efformantes.

Labia crassa: Opercula mutica, squamosa.

CHARACTER SPECIFICUS, &c.

Sparus flavus, pinnis virentibus, dorsali analique subfalcatis.

Gen. Zool. 4. p. 409.

Sparus falcatus. S. spinis quatuor in pinna ani.

Bloch. t. 258.

Elegans hæc species maria incolit Americana, ad longitudinem pertingens plus bipedalem. Primus eam descripsisse et depinxisse videtur celeberrimus Plumierus.

807



R. Nodder. Del. & Sculp.

THE
FALCATED SPARUS.

GENERIC CHARACTER.

Teeth strong: Front-teeth in some species disposed in a single row, in others in a double, treble, or quadruple row.

Grinders (in most species) convex, smooth, and disposed in ranges, forming a kind of pavement in the mouth.

Lips thick: *Gill-covers* unarmed, smooth, scaly.

SPECIFIC CHARACTER, &c.

Yellow Sparus with greenish fins; the dorsal and anal fin subfalcated.

Falcated Sparus.

Gen. Zool. 4. p. 409.

This elegant species is a native of the American seas, where it arrives at the length of more than two feet. It seems to have been first described and figured by the celebrated Plumier.

SCARABÆUS TYPHON.

CHARACTER GENERICUS.

Antennæ clavatæ capitulo fissili.

Tibiæ anticæ sæpius dentatæ.

Lin. Syst. Nat.

CHARACTER SPECIFICUS, &c.

Scarabæus scutellatus niger, thorace tricorni,
intermedio emarginato, capitis cornu erecto
recurvo bifido.

Scarabæus Typhon.

Olivier coleopt. Scarab. p. 12.
pl. 16. f. 152.

Indiam incolit Scarabæus Typhon, magnitudine
vera in tabula depictus.



Stag Beetle

TYPHON.

GENERIC CHARACTER.

Antennæ clavated with a fissile or laminated tip.

Middle joints of the fore-legs toothed.

SPECIFIC CHARACTER, &c.

Black scutellated Beetle, with three-horned thorax, the intermediate one emarginated, and an upright recurved bifid horn on the head.

Typhon.

Oliv. ins. coleopt. Scarab. p. 12.
pl. 16. f. 152.

The *Scarabæus Typhon* is a native of India, and is represented on the plate in its natural size.

BUCEROS ALBIROSTRIS.

CHARACTER GENERICUS.

Rostrum convexum, curvatum, cultratum, magnum, extrorsum serratum : frontis calvaria nuda, osseo-gibbosa.

Nares pone rostri basin.

Lingua acuta, brevis.

Pedes gressorii.

Lin. Syst. Nat.

CHARACTER SPECIFICUS, &c.

Buceros subcristatus niger, abdomine albo, fronte ossea antrorsum macula nigra.

Le Calao à bec blanc.

Levaill. ois. nouv. pl. 14.

Speciem hanc, Indiæ incolam, novam esse nec antea descriptam opinatur Levaillant. Ab apice rostri ad extremum caudæ longa est quasi viginti uncias. De modo vivendi nihil compertum est.



THE
WHITE-BEAKED HORNBILL.

GENERIC CHARACTER.

Bill very large, convex, curved, with a large bony frontal prominence: mandibles serrated, the serratures pointing outward.

Nostrils behind the base of the bill.

Tongue sharp, short.

Feet gressorial.

SPECIFIC CHARACTER, &c.

Subcristated black Hornbill, with white abdomen, and bony front with a black spot on the fore-part.

Le Calao à bec blanc.

Levaill. ois. nouv.

This bird Mons. Levaillant considers as a new species of the genus *Buceros*. It is a native of India, and measures about twenty inches from the tip of the bill to the end of the tail. Its manners and particular history are at present unknown.

SPARUS VERRES.

CHARACTER GENERICUS.

Dentes validi: primores in aliis ordine simplici, in aliis duplici, triplici, vel quadruplici dispositi.

Molares (plerisque) validi, convexi, læves, in series dispositi, et quasi pavementum in ore efformantes.

Labia crassa: opercula mutica, squamosa.

CHARACTER SPECIFICUS, &c.

Sparus albidus, dorso violaceo, pinna caudali rubra, dorsali et anali apice rubris.

Lutianus Verres.

Bloch ichth. t. 255.

Docet Blochius in mari Japonico generari Sparum Verrem, dentibus primoribus magnis et curvatis in utraque maxilla sitis notabilem. In magnitudinem, ut plurimum, crescit decem seu duodecim unciarum. Cuinam vivendi modo se præcipue assuescat, incertum est.

870



Thalassoma kribia

THE
BOAR-TUSKED SPARUS.

GENERIC CHARACTER.

Teeth strong: front teeth in some species disposed in a single row, in others in a double, treble, or quadruple row.

Grinders (in most species) convex, smooth, and disposed in ranges, forming a kind of pavement in the mouth.

Lips thick: gill-covers unarmed, smooth, scaly.

SPECIFIC CHARACTER, &c.

Whitish Sparus, with the back violet, the tail and tips of the dorsal and anal fin red.

Boar Sparus.

Gen. Zool. 4. p. 414.

This fish is, according to Dr. Bloch, a native of the Japanese sea, and is remarkable for the large size and curvature of the front teeth in both jaws. Its general length is about ten or twelve inches, but its particular history seems to be in a great degree unknown.

LUCANUS ALCES.

CHARACTER GENERICUS.

Antennæ clavatæ; clava compressa latere latiore
pectinato-fissili.

Maxillæ porrectæ, exsertæ, dentatæ.

Lin. Syst. Nat.

CHARACTER SPECIFICUS, &c.

Lucanus niger, maxillis exsertis dente medio
validissimo, apice quadridentatis.

Lucanus Alces. *L. mandibulis* exsertis apice
quadridentatis.

Fabr. sp. ins. p. 1.

Olivier. entom. 1. p. 8.

Petiv. gazoph. t. 47. f. 15.

caput.

Indiam incolit rara hæc Lucani species, et magni-
tudine vera in tabula depingitur.

877



R. P. W. Decker Del. et Sculp.

THE
ELK-HORNED STAG-CHAFFER.

GENERIC CHARACTER.

Antennæ clavated, with compressed tip, divided into lamellæ on the inner side.

Jaws stretched forwards, exserted, and toothed.

SPECIFIC CHARACTER, &c.

Black Stag-Chaffer, with exserted jaws, furnished on the middle with a very strong tooth, and four-toothed at the tips.

The Elk-Horned Lucanus.

Black Indian Stag-Chaffer.

This rare species of Lucanus is a native of India, and is represented on the plate in its natural size.

VOLUTA MAGNIFICA.

CHARACTER GENERICUS.

Animal Limax.

Testa unilocularis, spiralis.

Apertura ecaudata, subeffusa.

Columella plicata, Labio Umbilicove nullo.

Lin. Syst. Nat.

CHARACTER SPECIFICUS, &c.

Voluta ventricosa trifasciata, albo fuscoque transversim varia, columella quadriplicata

Voluta magnifica. V. testa ovali glabra ventricosa tumida, fasciis latis circumcincta, maculis fasciarum saturatioribus, anfractibus sex, apertura lunari amplissima ex croceo et aurantio infecta, apice obtuso papillari, basi emarginata, columella quadriplicata.

Chemn. 11. p. 8. t. 174, 175.

E plurimis speciebus quas continet genus Voluta, rarior et elegantior nulla est illa quam depinximus; quæ in oceano pacifico australi generata, in Angliam illata est quo tempore circa orbem terrarum primum navigabat celeberrimus Cookius. In tabula nostra magnitudine circiter tertia parte deminuta exprimitur.

THE
MAGNIFIC VOLUTE.

GENERIC CHARACTER.

Animal resembling a Limax or Slug.

Shell unilocular, spiral.

Aperture somewhat spreading, simple.

Pillar wreathed or pleated.

SPECIFIC CHARACTER, &c.

Trifasciated ventricose Volute, with brown and white transverse variegations, and three-pleated pillar.

The magnificent Volute.

Great Southern variegated Volute.

The present shell may justly be numbered among the rarest and most elegant species of the numerous genus to which it belongs. It is a native of the Southern Pacific Ocean, and was brought into this country by the naturalists who accompanied Captain Cook in his first circumnavigation. The plate represents it about a third less than the natural size.

AMPELIS POMPADORA.

CHARACTER GENERICUS.

Rostrum rectum, convexum ; mandibula superiore longiore, subincurvata, utrinque emarginata.

Lingua acuta, cartilaginea, bifida.

Lin. Syst. Nat.

CHARACTER SPECIFICUS, &c.

Ampelis Pompadora. A. purpurea, tectricibus alarum proximis ensiformibus elongatis carinatis nudis.

Lin. Syst. Nat.

Cotinga purpurea.

Briss. av. 2. p. 347. t. 35. f. 1.

Turdus puniceus.

Pall. adumbr. 99.

Formosissimam hanc avem a Brissonio primitus descriptam et depictam, generat America australis, paulo majorem Garrulo Linnæi seu Ampelide Europea. Testatur Vaillantus feminam a mari in hoc differre, quod pars superior multo magis infussetur, cum inferior alba sit, purpureo suffusa.



22. Not. 2. d. 1. 1. 1.

THE
POMPADOUR CHATTERER.

GENERIC CHARACTER.

Bill strait, convex; the upper mandible rather longer, subincurvated, emarginated on each side near the tip.

Tongue sharp-pointed; cartilaginous, bifid.

SPECIFIC CHARACTER, &c.

Purple Chatterer, with white quill-feathers; the coverts nearest the body long, narrow, stiff, and ensiform.

The Pompadour Chatterer.

Edwards, pl. 341.

Le Cotinga Pacapaca.

Levaill. ois. amer. pl. 31.

This highly beautiful bird is a native of South America, and seems to have been first described and figured by the accurate Brisson. In size it is rather superior to the *Ampelis Garrulus* of Linnæus, or common European Chatterer. The female, according to Mons. Le Vaillant, differs in being of a much browner colour on the upper parts, and white, slightly tinged with purple, beneath.

EXOCOETUS MESOGASTER.

CHARACTER GENERICUS.

Caput squamosum: maxillis utroque latere connexis.

Membr: branch: radiis decem.

Pinnæ pectorales maximæ, volatiles.

CHARACTER SPECIFICUS, &c.

Exocoetus argenteo-cærulescens, pinnis ventralibus in medio abdominis.

Gen. Zool. 5. p. 146.

Exocoetus Mesogaster. E. pinnis ventralibus in æquilibrio.

Bloch. ichth. t. 399.

A congeneribus differt *Exocoetus Mesogaster* situ pinnarum ventralium. In maribus Americanis generatur, et ad longitudinem pertingit pedalem, vel etiam sesquipedalem.

814



Pezomyxus volitans

THE
MIDDLE-FINNED FLYING-FISH.

GENERIC CHARACTER.

Head scaly : jaws connected on each side.

Gill-Membrane ten-rayed.

Pectoral fins very large, giving the power of flight.

SPECIFIC CHARACTER, &c.

Silvery-blueish Flying-Fish, with the ventral fins situated on the middle of the abdomen.

American Flying-Fish.

Gen. Zool. 5. p. 146.

The *Exocoetus Mesogaster* differs from the rest of its congeners in the situation of the ventral fins. It is a native of the American seas, and grows to the length of twelve or eighteen inches.

VOLUTA JUNONIA.

CHARACTER GENERICUS.

Animal Limax.

Testa unilocularis, spiralis.

Apertura ecaudata, subeffusa.

Columella plicata.

Lin. Syst. Nat.

CHARACTER SPECIFICUS, &c.

Voluta subventricosa flavescens, maculis subquadratis transversis rufis, spira longitudinaliter striata, columella septemplicata.

Voluta Junonia. V. rarissima ovata lævis, maculis quadratis rufescentibus in fundo flavido seriatim stratis nitidissime tessulata et zonata, spira exserta striata, apice semigloboso, basi transversim striata, subemarginata, columella septemplicata.

Chemn. 11. p. 16. t. 177.

Rarissimam hanc speciem, in maribus, ut creditur, australibus innatam, jactant perpauca Musea Europea. Primus eam descripsisse videtur Dominus Favanne in opere conchyliologico Domini Argenville, quod auctius edidit. Tabula exprimit magnitudinem naturalem.



THE
JUNONIAN VOLUTE.

GENERIC CHARACTER.

Animal resembling a Limax or Slug.

Shell unilocular, spiral.

Aperture somewhat spreading, simple.

Pillar wreathed or pleated.

SPECIFIC CHARACTER, &c.

Yellowish subventricose Volute, with squarish transverse rufous spots, longitudinally striated spire, and seven-pleated pillar.

Voluta Junonia.

Chemn. 11. p. 16. pl. 177.

Favanne Argenv.

This extremely rare species, which makes its appearance in but very few of the European cabinets, is supposed to be a native of the Southern Pacific, and seems to have been first described by Mons. Favanne, in his enlarged edition of Argenville's Conchyliology. It is represented on the plate in its natural size.

PHALÆNA COSSUS.

CHARACTER GENERICUS.

Antennæ setaceæ, a basi ad apicem sensim
attennatæ.

Alæ (sedentis) sæpius deflexæ, (volatu nocturno).

Lin. Syst. Nat.

CHARACTER SPECIFICUS, &c.

Phalæna cinerea nigro reticulata, alis deflexis,
thorace postice fascia atra.

Phalæna Cossus. P. (*Bombyx*) elinguis, alis
deflexis nebulosis, thorace postice fascia,
atra, antennis lamellatis.

Lin. Syst. Nat.

In maximis speciebus Britannicis numeranda *Phalæna Cossus* e larva exoritur cui facies nec jucunda nec elegans; quæque salictis, quorum in truncis præcipue delitescit, lignum exedendo pestifera est et exitiosa. Odorem præterea spargit teterrimum, quo arbores in quibus sibi sedem elegerit, dignosci possint. Quum ad tertium ætatis annum pervenerit larva, creditur in chrysalidem converti, plerumque mense Maio, postquam se tela seu theca valida e parvulis fibris ligneis suo ipsius glutine conjunctis involverit, cortici ut plurimum arboris inferiori affixa; unde mense Junio e

carcere sese liberat phalæna, e theca nempe protrusa chrysalide ope spiculorum fortium quibus annuli corporis armantur.

Nefas esset de hoc insecto disserenti silentio præterire eximium Lyonetti opus, qui de dissectione larvæ locutus, plures in illa numerat musculos quam in ipso humano corpore. Pari diligentia in singulis ipsius phalænæ partes anatomice inquisivit idem ingeniosus auctor.



THE
GOAT MOTH.

GENERIC CHARACTER.

Antennæ setaceous, gradually lessening from the base to the tip.

Wings (when sitting) generally deflected. Flight diurnal.

SPECIFIC CHARACTER, &c.

Grey Moth, with black reticulations, deflected wings, and a black stripe across the hind part of the thorax.

Great Willow-wood Moth.

Goat Moth.

Albin. ins. pl. 35.

Roes. ins. 1. *Phal.* Cl. 2. pl. 18.

The *Phalæna Cossus* or Goat Moth, one of the largest of the British *Phalænæ*, proceeds from a larva or caterpillar of an appearance rather disagreeable than elegant. It resides in the trunks of trees, more particularly of willows; feeding on the wood, and committing great ravages in districts where it abounds. It diffuses a highly unpleasing scent, by which its residence may frequently be known in passing by such trees as are much infested by it. It is generally sup-

posed to arrive at its third year before it changes to a chrysalis, which usually takes place in the month of May, the animal first enveloping itself in a strong web or covering, formed of particles of wood united by a strong gluten: this web is usually affixed to the bark of the lower part of the tree; and from it, in the month of June, proceeds the complete insect, the chrysalis forcing its way through the web by the assistance of the strong spiculæ with which its segments are surrounded.

It would be unpardonable, in giving the history of this insect, to omit mentioning the celebrated publication of Lyonett, relative to the anatomy of its caterpillar, which, according to that author, is furnished with a far greater number of muscles than the human body itself. The same writer afterwards pursued his researches with equal diligence into the anatomy of the complete insect.

AMPELIS COTINGA.

CHARACTER GENERICUS.

Rostrum rectum, convexum; mandibula superiore longiore, subincurvata, utrinque emarginata.

Lingua acuta, cartilaginea, bifida.

Lin. Syst. Nat.

CHARACTER SPECIFICUS, &c.

Ampelis Cotinga. A. nitidissima cærulea, subtus purpurea, alis caudaque nigris.

Lin. Syst. Nat.

Cotinga.

Brisson av. 2. p. 340.

Omnium congenerum splendidissima est *Ampelis Cotinga*, eadem magnitudine qua *Ampelis Pompadora*: easdem etiam regiones Americanas incolit. Coloribus interdum insigniter variat.



THE
PURPLE-BREASTED CHATTERER.

GENERIC CHARACTER.

Bill strait, convex; upper mandible rather longer, subincurvated, emarginated on each side near the tip.

Tongue sharp-pointed, cartilaginous, bifid.

SPECIFIC CHARACTER, &c.

Brilliant blue Chatterer, purple beneath, with black wings and tail.

Le Cotinga du Bresil.

Pl. Enl. 186.

Purple-Breasted Manakin.

Edw. pl. 241.

This, which is the most splendid of the whole genus to which it belongs, is of similar size with the *Ampelis Pompadora*, and inhabits the same regions. In colour it occasionally varies considerably.

9/8



R. N. S. D. & S. S. S. S.

THE
SURINAM SPARUS.

GENERIC CHARACTER.

Teeth strong: Front-teeth in some species disposed in a single row, in others in a double, treble, or quadruple row.

Grinders (in most species) convex, smooth, and disposed in ranges, forming a kind of pavement in the mouth.

Lips thick: *Gill-covers* unarmed, smooth, scaly.

SPECIFIC CHARACTER, &c.

Reddish Spare, with black semidecurrent bands, and whitish abdomen with a black spot on each side.

Der Steinkahlkopf.

Bloch ichth. pl. 253.

Surinam Sparus.

Gen. Zool. 4. p. 458.

This fish is a native of the American seas, measuring about ten or twelve inches in length.

PAPILIO PIERA.

CHARACTER GENERICUS.

Antennæ apicem versus crassiores, sæpius clavato-capitatae.

Alæ (sedentis) erectæ sursumque conniventes.
(Volatu diurno.)

Lin. Syst. Nat. p. 744.

CHARACTER SPECIFICUS, &c.

Papilio Piera. P. alis oblongis integerrimis
hyalinis diaphanis, posticis ocellis binis.

Lin. Syst. Nat.

Mer. Sur. t. 16.

Roes. Append. t. 6.

Surinamiam incolit Papilio Piera, ubi larva ejus,
testante Domina Merian, folia Anacardii occidentalis
depascitur.



THE
TALC-WINGED BUTTERFLY.

GENERIC CHARACTER.

Antennæ commonly thickening towards the end
into a clavated tip.

Wings (when at rest) meeting upwards. Flight
diurnal.

SPECIFIC CHARACTER, &c.

Butterfly with oblong, entire, transparent
wings; the lower pair marked by two
ocellated spots.

Mer. Sur. pl. 16.

Roes. Append. pl. 6.

This insect is a native of Surinam, where its larva
or caterpillar, according to Madam Merian, feeds on
the leaves of the *Anacardium occidentale* or American
Cashew.

MADREPORA AXILLARIS.

CHARACTER GENERICUS.

Animal Medusa.

Corallium cavitatibus lamelloso-stellatis.

Lin. Syst. Nat.

CHARACTER SPECIFICUS.

Madrepora dichotoma, ramis distinctis divaricatis, stellis terminalibus turbinatis; axillaribus compressis; centris dilatatis exesis.

Soland. et Ellis Zooph. p. 153.

t. 13. f. 5.

Maria incolit Indica Madrepora axillaris, et magnitudine vera in tabula depingitur.



29. Noddy Del et Sculp.

THE
AXILLARY MADREPORE.

GENERIC CHARACTER.

Animal allied to a Medusa.

Coral marked by radiating lamellar cavities.

SPECIFIC CHARACTER, &c.

Dichotomous Madrepore, with distinct divaricated branches; the axillary stars compressed; the terminal ones turbinated; with dilated and excavated centres.

Madrepora axillaris.

Soland. et Ellis Zooph. p. 153.
pl. 13. f. 5.

The present species of Madrepore is an inhabitant of the Indian seas, and is represented on the plate in its natural size.



THE
SUPERB CHATTERER.

GENERIC CHARACTER.

Bill strait, convex: upper mandible rather longer, subincurvated, emarginated on each side near the tip.

Tongue cartilaginous, sharp-pointed, bifid.

SPECIFIC CHARACTER; &c.

Brilliant blue Chatterer, deep purple beneath, varied with fulvous-crimson spots.

Purple-Breasted Chatterer. *Var.?*

The highly elegant and rare bird represented on the present plate, has been generally considered as the male of the *Ampelis Cotinga*, or Purple-Breasted Chatterer. Mons. Levaillant, however, so well known for his extensive information, is inclined to consider it as constituting at least a remarkable variety, if not a distinct species. It inhabits the same regions with the former kind.

PHALÆNA PINI.

CHARACTER GENERICUS.

Antennæ setaceæ, a basi ad apicem sensim attenuatæ.

Alæ (sedentis) sæpius deflexæ (volatu diurno).

Lin. Syst. Nat.

CHARACTER SPECIFICUS, &c.

Phalæna (*Bombyx*) *elinguis*, alis reversis griseis, strigis duabus cinereis, puncto albo triangulari.

Lin. Syst. Nat. p. 814.

Merian Ins. Europ. t. 22.

Frisch Ins. 10. t. 10.

Rarior longe in Britannia quam in reliqua omni Europa *Phalæna Pini* folia depascitur pinorum et abietum. Mense Junio, postquam tela sese tenui et quasi ovata involverit, in chrysalidem convertitur; e qua insequente Julio erumpit insectum plene formatum.



R. P. Nodder. Sculp^t.

THE
PINE MOTH.

GENERIC CHARACTER.

Antennæ setaceous, gradually lessening from base to tip.

Wings (when sitting) often deflex. Flight diurnal.

SPECIFIC CHARACTER, &c.

Tongueless Moth, with reversed grey wings, with two cinereous streaks, and a triangular white speck.

Merian. Eur. Ins. pl. 22.

Roes. Ins. 1. pl. 59.

This insect, which is much more common in other European regions than in our own country, feeds, during its caterpillar state, on the leaves of Pine and Fir trees, and changes into a chrysalis in the month of June, first enveloping itself in a loose oval web; and the Moth makes its appearance in July.

PLEURONECTES ARGUS.

CHARACTER GENERICUS.

Oculi ambo in eodem latere capitis.

Corpus compressum, latere altero dorsum, altero abdomen referente.

CHARACTER SPECIFICUS, &c.

Pleuronectes albidus, maculis rotundis flavis
cæruleo marginatis.

Pleuronectes Argus. P. corpore vario, pinna
caudæ rotundata,

Bloch. t. 48.

Passer oculatus.

Plumier. Mss.

Maria incolit Americana *Pleuronectes Argus*, in
longitudinem pedalem vel sesquipedalem crescens.



THE
ARGUS FLOUNDER.

GENERIC CHARACTER.

Eyes both on the same side of the head.

Body compressed; one side representing the back, the other the abdomen.

SPECIFIC CHARACTER, &c.

Whitish Flounder, with round yellow spots margined with blue.

Argus Flounder.

Gen. Zool. 4. p. 319.

The Argus Flounder is an inhabitant of the American seas, and grows to the length of twelve or eighteen inches.

PAPILIO RHAMNÍ.

CHARACTER GENERICUS.

Antennæ apicem versus crassiores, sæpius clavato-capitatae.

Alæ (seditis) erectæ sursumque conniventes. (Volatu diurno).

Lin. Syst. Nat.

CHARACTER SPECIFICUS, &c.

Papilio Rhamni. P. alis integerrimis angulatis flavis: singulis puncto fulvo, subtus, ferrugineo.

Lin. Syst. Nat.

Degeer. Ins. 1. t. 15.

Roes. Ins. 3. t. 46.

Folia Rhamni cathartici præcipue depascitur larva hujus papilionis, et in chrysalidem mense Maio convertitur, e qua mense Junio erumpit Papilio.

324



THE
BUCKTHORN BUTTERFLY.

GENERIC CHARACTER.

Antennæ commonly thickening towards the end
into a clavated tip.

Wings (when at rest) meeting upwards. Flight
diurnal.

SPECIFIC CHARACTER, &c.

Butterfly with yellow, entire, angulated wings,
with a fulvous spot on each, beneath fer-
ruginous.

The Buckthorn Butterfly.

The Brimstone Butterfly.

The caterpillar of elegant insect represented on the
present plate, feeds principally on the leaves of Buck-
thorn, changing into a chrysalis in May, and giving
birth to the Butterfly in June.

BODIANUS AYA.

CHARACTER GENERICUS.

Habitus Percæ.

Opercula squamosa, aculeata.

Squamæ (pluribus) læves.

CHARACTER SPECIFICUS, &c.

Bodianus ruber, abdomine argenteo, operculis unispinosus, cauda lunata.

Gen. Zool. 4. p. 570.

Bodianus Aya. *B. aculeo* ad operculum, pinna caudæ lunata.

Bloch. ichth. 7. t. 227.

Lacus incolit Brasilienses *Bodianus Aya*, inter congeneres forte puererrimus. In longitudinem tripedalem crescit, et in cibis habetur delicatioribus.

822



Thalassoma daniellii

THE
AYA BODIAN.

GENERIC CHARACTER.

Habit of the genus *Perca*.

Gill-Covers aculeated.

Scales (in most species) smooth.

SPECIFIC CHARACTER, &c.

Red Bodian, with silvery abdomen, single-spined gill-covers, and lunated tail.

Bodianus Aya.

Bloch. ichth. pl. 227.

Acara Aya.

Will. ichth. p. 329.

pl. x. 7. f. 6.

The fish here represented is an inhabitant of the Brazilian lakes, and is, perhaps, the most beautiful of its congeners. It grows to the length of three feet, and is considered as a very delicate article of food.

VOLUTA INDICA.

CHARACTER GENERICUS.

Animal Limax.

Testa unilocularis, spiralis.

Apertura ecaudata, subeffusa.

Columella plicata.

Lin. Syst. Nat.

CHARACTER SPECIFICUS, &c.

Voluta Indica. V. testa elongata flava, fasciis
tribus macularum fuscarum, columella
quadriplicata.

Lin. Gmel. p. 3467.

Cymbium basi coarctata luteum, quadruplici
ordine macularum nigrescentium fascia-
tum, quadriplicatum.

Martini conch. 3. t. 72. f. 772. 773.

In maribus Indicis generatur rara hæc Volutæ spe-
cies, et magnitudine vera in tabula exprimitur.

827



R.P.N.

THE
INDIAN VOLUTE.

GENERIC CHARACTER.

Animal resembling a Limax or Slug.

Shell unilocular, spiral.

Aperture somewhat spreading, simple.

Pillar wreathed or pleated.

SPECIFIC CHARACTER, &c.

Yellowish oblong Volute, with three bands of irregular brown spots, and four-pleated pillar.

Couronne ethiopique mouchetée.

Knorr. Vergn. 5. pl. 8. f. 1.

The Spotted Indian Volute.

This rare species of Volute is an inhabitant of the Indian seas, and is figured on the annexed plate in its natural size.

PAPILIO EMPEDOCLES.

CHARACTER GENERICUS.

Antennæ apicem versus crassiores, sæpius clavato-capitatae.

Alæ (sedentis) erectæ sursumque conniventes, (volatu diurno.)

Lin. Syst. Nat.

CHARACTER SPECIFICUS, &c.

Papilio antennis setaceis, alis fuscis luteo striatis stria media latiore, posticis ocellato-caudatis.

Papilio *Ægisthus*. *P.* alis fusco luteoque undulatis, ocellis caudalibus subternis.

Lin. Gmel. p. 2240.

Phalæna *Lunus*. *P.* (A.) seticornis spirilignis, alis caudatis fusco luteoque undulatis, ocellis caudalibus subternis.

Lin. Syst. Nat. p. 810.

Papilio *Empedocles*.

Cramer. t. 199.

Papilio *Lunus*.

Cramer. t. 200.?

Surinamiam incolit *Papilio* *Empedocles*, et magnitudine vera in tabula depingitur.

EMPEDOCLES.

GENERIC CHARACTER.

Antennæ thickening towards the end, and commonly terminating in a clavated tip.

Wings (when at rest) meeting upwards. (Flight diurnal.)

SPECIFIC CHARACTER, &c.

Butterfly with setaceous antennæ, brown wings with luteous streaks, the middle streak broadest; the lower wings eye-tailed.

Papilio Empedocles.

Cram. t. 199.

Papilio Ægisthus.

Lin. Gmel.

Phalæna Lunus.

Lin. Syst. Nat.

This insect is a native of Surinam, and is represented on the plate in its natural size.



INDEX.

FL.

- 802. *Acanthurus Chirurgus*.
- 793. *Alcedo leucocephala*.
- 825. — *Smyrnensis*.
- 813. *Ampelis Pompadora*.
- 817. — *Cotinga*.
- 821. — *superba*.
- 792. *Aphrodita cirrhosa*.
- 796. *Brachionus aculeatus*.
- 826. *Bodianus Aya*.
- 799. *Buccinum subulatum*.
- 809. *Buceros albirostris*.
- 797. *Certhia gutturalis*.
- 782. *Cephalus brevis*.
- 800. *Chætodon Teira*.
- 781. *Coracias varia*.
- 785. *Columba aromatica*.
- 814. *Exocoëtus mesogaster*.
- 790. *Fistularia Chinensis*.
- 784. *Gryllus nasutus*.
- 788. — *squarrosus*.
- 786. *Labrus trimaculatus*.
- 811. *Lucanus Alces*.
- 820. *Madrepore axillaris*.
- 783. *Papilio Adippe*.
- 787. — *Amphinome*.
- 791. — *Cassia*.
- 804. — *Androgeus*.
- 806. — *Spondiæ*.
- 819. — *Piera*.
- 824. — *Rhamni*.
- 828. — *Empedocles*.
- 816. *Phalæna Cossus*.
- 822. — *Pini*.
- 823. *Pleuronectes Argus*.
- 769. *Psittacus infuscatus*.
- 808. *Scarabæus Typhon*.
- 794. *Scarus Cretensis*.
- 807. *Sparus falcatus*.
- 810. — *Verres*.
- 818. — *Surinamensis*.
- 803. *Sphinx populi*.
- 795. — *satellitæ*.
- 801. *Strix larvata*.
- 805. *Sturnus subroseus*.
- 798. *Tubularia ramosa*.
- 812. *Voluta magnifica*.
- 815. — *Junonia*.
- 827. — *Indica*.

PL.

- 802. *Acanthurus lancet-tailed*.
- 792. *Aphrodita cirrhated*.
- 808. *Beetle Typhon*.
- 826. *Bodian Aya*.
- 796. *Brachionus aculeated*.
- 799. *Buccinum subulate*.
- 783. *Butterfly Adippe*.
- 787. — *Amphinome*.
- 791. — *Cassia*.
- 804. — *Androgeus*.
- 806. — *single-spotted*.
- 819. — *talc-winged*.
- 824. — *buckthorn*.
- 828. — *Empedocles*.
- 800. *Chætodon long-finned*.
- 797. *Chatterer Pompadour*.
- 817. — *purple-breasted*.
- 821. — *superb*.
- 797. *Creeper purple-throated*.
- 790. *Fistularia Chinese*.
- 823. *Flounder Argus*.
- 814. *Flying-Fish middle-finned*.
- 809. *Hornbill white-beaked*.
- 825. *Kingfisher Smyrna*.
- 793. — *white-headed*.
- 786. *Labrus trimaculated*.
- 784. *Locust long-fronted*.
- 788. — *triple-spined*.
- 820. *Madrepore axillary*.
- 816. *Moth Goat*.
- 822. — *pine*.
- 801. *Owl masked*.
- 789. *Parrot dusky*.
- 785. *Pigeon aromatic*.
- 781. *Roller pied*.
- 794. *Scarus Cretan*.
- 810. *Sparus boar-tusked*.
- 807. — *falcated*.
- 818. — *Surinam*.
- 795. *Sphinx double-pointed*.
- 803. — *poplar*.
- 782. *Sun-Fish short*.
- 811. *Stag-Chaffer elk-horned*.
- 805. *Stare subrosaceous*.
- 798. *Tubularia twisted*.
- 812. *Volute magnific*.
- 815. — *Junonian*.
- 827. — *Indian*.

Acme Library Card Pocket

Under Pat. Sept. 26, '76, "Ref. Index File"

Made by **LIBRARY BUREAU**

CHICAGO. NEW YORK. BOSTON.

Keep Your Card In This Pocket

